

D-Glucose

UV-method

for the determination of D-glucose in foodstuffs and other materials

Cat. No. 10 716 251 035

Test-Combination for 3 × 45 determinations

BOEHRINGER MANNHEIM / R-BIOPHARM
Enzymatic BioAnalysis / Food Analysis


For use in *in vitro* only

Store at 2-8°C


For recommendations for methods and standardized procedures see references (A 2, B 2, C 2, D 2)

Principle (Ref. A 1)

D-Glucose is phosphorylated to D-glucose-6-phosphate (G-6-P) in the presence of the enzyme hexokinase (HK) and adenosine-5'-triphosphate (ATP) with the simultaneous formation of adenosine-5'-diphosphate (ADP) (1).



In the presence of the enzyme glucose-6-phosphate dehydrogenase (G6P-DH), G-6-P is oxidized by nicotinamide-adenine dinucleotide phosphate (NADP) to D-gluconate-6-phosphate with the formation of reduced nicotinamide-adenine dinucleotide phosphate (NADPH) (2).



The amount of NADPH formed in this reaction is stoichiometric to the amount of D-glucose. The increase in NADPH is measured by means of its light absorbance at 334, 340 or 365 nm.

The Test-Combination contains

- Three bottles 1, each with approx. 7.2 g powder mixture, consisting of: triethanolamine buffer, pH approx. 7.6; NADP, approx. 110 mg; ATP, approx. 260 mg; magnesium sulfate
- Three bottles 2, each with approx. 1.1 ml suspension, consisting of: hexokinase, approx. 320 U; glucose-6-phosphate dehydrogenase, approx. 160 U
- Bottle 3 with D-glucose assay control solution for assay control purposes (measurement of the assay control solution is not necessary for calculating the results.) Use the assay control solution undiluted. (Expiry date: see pack label)

Preparation of solutions

- Dissolve contents of one bottle 1 with 45 ml redist. water.
- Use contents of one bottle 2 undiluted.

Stability of reagents

The contents of the bottles 1 are stable at 2-8°C (see pack label).

Solution 1 is stable for 4 weeks at 2-8°C, and for 2 months at -15 to -25°C.

Bring solution 1 to 20-25°C before use.

The contents of the bottles 2 are stable at 2-8°C (see pack label).

Procedure

Wavelength¹: 340 nm, Hg 365 nm or Hg 334 nm

Glass cuvette²: 1.00 cm light path

Temperature: 20-25°C

Final volume: 3.020 ml

Read against air (without a cuvette in the light path) or against water

Sample solution: 1-100 µg of D-glucose/assay³ (in 0.100-2.000 ml sample volume).

Pipette into cuvettes	Blank	Sample
solution 1	1.000 ml	1.000 ml
sample solution*	-	0.100 ml
redist. water	2.000 ml	1.900 ml
Mix** and read absorbances of the solutions (A ₁) after approx. 3 min. Start reaction by addition of:		
suspension 2	0.020 ml	0.020 ml
Mix**, wait until the reaction has stopped (approx. 10-15 min) and read the absorbances of the solutions (A ₂). If the reaction has not stopped after 15 min, continue to read the absorbances at 2 min intervals until the absorbance increases constantly for 2 min.***		

* Rinse the enzyme pipette or the pipette tip of the piston pipette with sample solution before dispensing the sample solution.

** For example, with a plastic spatula or by gentle swirling after closing the cuvette with Parafilm (trademark of the American Can Company, Greenwich, Ct., USA)

*** Creep reactions occur very occasionally. They are mostly brought about by the contents of the sample solutions, such as enzymes or coloring agents. These interfering substances may be removed during sample preparation.

If the absorbance A₂ increases constantly, extrapolate the absorbance to the time of the addition of suspension 2 (HK/G6P-DH).

Determine the absorbance differences (A₂-A₁) for both, blank and sample. Subtract the absorbance difference of the blank from the absorbance difference of the sample.

$$\Delta A = (A_2 - A_1)_{\text{sample}} - (A_2 - A_1)_{\text{blank}}$$

The measured absorbance differences should, as a rule, be at least 0.100 absorbance units to achieve sufficiently precise results (see "Instructions for performance of assay" and "Sensitivity and detection limit", pt. 3).

Calculation

According to the general equation for calculating the concentration:

$$c = \frac{V \times MW}{\epsilon \times d \times v \times 1000} \times \Delta A \text{ [g/l]}$$

V = final volume [ml]

v = sample volume [ml]

MW = molecular weight of the substance to be assayed [g/mol]

d = light path [cm]

ε = extinction coefficient of NADPH at

$$340 \text{ nm} = 6.3 \text{ [l} \times \text{mmol}^{-1} \times \text{cm}^{-1}\text{]}$$

$$\text{Hg } 365 \text{ nm} = 3.5 \text{ [l} \times \text{mmol}^{-1} \times \text{cm}^{-1}\text{]}$$

$$\text{Hg } 334 \text{ nm} = 6.18 \text{ [l} \times \text{mmol}^{-1} \times \text{cm}^{-1}\text{]}$$

It follows for D-glucose:

$$c = \frac{3.020 \times 180.16}{\epsilon \times 1.00 \times 0.100 \times 1000} \times \Delta A_{\text{D-glucose}} = \frac{5.441}{\epsilon} \times \frac{\Delta A_{\text{D-Glucose}}}{\text{[g D-glucose/l sample solution]}}$$

If the sample has been diluted during preparation, the result must be multiplied by the dilution factor F.

When analyzing solid and semi-solid samples which are weighed out for sample preparation, the result is to be calculated from the amount weighed.

$$\text{Content}_{\text{D-glucose}} = \frac{c_{\text{D-glucose}} \text{ [g/l sample solution]}}{\text{weight}_{\text{sample}} \text{ in g/l sample solution}} \times 100 \text{ [g/100 g]}$$

1. Instructions for performance of assay

The amount of D-glucose present in the assay has to be between 2 µg and 100 µg (measurement at 365 nm) or 1 µg and 50 µg (measurement at 340, 334 nm), respectively. In order to get a sufficient absorbance difference, the sample solution is diluted to yield a D-glucose concentration between 0.15 and 1.0 g/l or 0.08 and 0.5 g/l, respectively.

Dilution table

Estimated amount of D-glucose per liter		Dilution with water	Dilution factor F
measurement at			
340 or 334 nm	365 nm		
< 0.5 g	< 1.0 g	-	1
0.5-5.0 g	1.0-10.0 g	1 + 9	10
5.0-50 g	10.0-100 g	1 + 99	100
> 50 g	> 100 g	1 + 999	1000

1 The absorption maximum of NADPH is at 340 nm. On spectrophotometers, measurements are taken at the absorption maximum; if spectralline photometers equipped with a mercury vapor lamp are used, measurements are taken at a wavelength of 365 nm or 334 nm.

2 If desired, disposable cuvettes may be used instead of glass cuvettes.

3 See instructions for performance of assay.



0603.117482620®



If the measured absorbance difference (ΔA) is too low (e.g. < 0.100), the sample solution should be prepared again (weigh out more sample or dilute less strongly) or the sample volume to be pipetted into the cuvette can be increased up to 2.000 ml. The volume of water added must then be reduced so as to obtain the same final volume in the assays for sample and blank. The new sample volume v must be taken into account in the calculation

2. Specificity (Ref. A 1)

The method is specific for D-glucose.

In the analysis of commercial water-free D-glucose (molecular weight 180.16) and D-glucose monohydrate (molecular weight 198.17), results of $< 100\%$ have to be expected because the materials absorb moisture.

3. Sensitivity and detection limit

The smallest differentiating absorbance for the procedure is 0.005 absorbance units. This corresponds to a maximum sample volume $v = 2.000$ ml and measurement at 340 nm of a D-glucose concentration of 0.2 mg/l sample solution (if $v = 0.100$ ml, this corresponds to 4 mg/l sample solution).

The detection limit of 0.4 mg/l is derived from the absorbance difference of 0.010 (as measured at 340 nm) and a maximum sample volume $v = 2.000$ ml.

4. Linearity

Linearity of the determination exists from approx. $1 \mu\text{g}$ D-glucose/assay (0.4 mg D-glucose/sample solution; sample volume $v = 2.000$ ml) to $100 \mu\text{g}$ D-glucose/assay (1 g D-glucose/sample solution; sample volume $v = 0.100$ ml).

5. Precision

In a double determination using one sample solution, a difference of 0.005 to 0.010 absorbance units may occur. With a sample volume of $v = 0.100$ ml and measurement at 340 nm, this corresponds to a D-glucose concentration of approx. 4-8 mg/l. (If the sample is diluted during sample preparation, the result has to be multiplied by the dilution factor F . If the sample is weighed in for sample preparation, e.g. using 1 g sample/100 ml = 10 g/l, a difference of 0.04-0.08 g/100 g can be expected.)

The following data have been published in the literature:

D-glucose in blood: CV = 1.2 % (Ref. A 1.1)
CV = 1.8 % (Ref. A 1.2)

Diet beer:

D-glucose:

$x = 1.0 \text{ g}/100 \text{ ml}$ $r = 0.030 \text{ g}/100 \text{ ml}$ $s_{(r)} = \pm 0.011 \text{ g}/100 \text{ ml}$
 $R = 0.122 \text{ g}/100 \text{ ml}$ $s_{(R)} = \pm 0.043 \text{ g}/100 \text{ ml}$

For further data see references (Ref. A 2.4)

Fruit juice:

D-glucose: $r = 0.42 + 0.027 \times (c_{\text{D-glucose}} \text{ in g/l}) \text{ g/l}$
 $R = 1.00 + 0.042 \times (c_{\text{D-glucose}} \text{ in g/l}) \text{ g/l}$

D-fructose:

$r = 0.15 + 0.033 \times (c_{\text{D-fructose}} \text{ in g/l}) \text{ g/l}$
 $R = 1.05 + 0.045 \times (c_{\text{D-fructose}} \text{ in g/l}) \text{ g/l}$

For further data see references (Ref. B 2.9)

Wine:

$r = 0.056 \times x_i$
 $R = 0.12 + 0.076 x_i$

$x_i = \text{D-glucose resp. D-fructose content in g/l}$ (Ref. B2.17, 2.18)

Liquid whole egg:

D-glucose:

$x = 0.44 \text{ g}/100 \text{ g}$ $r = 0.073 \text{ g}/100 \text{ g}$ $s_{(r)} = \pm 0.026 \text{ g}/100 \text{ g}$
 $R = 0.106 \text{ g}/100 \text{ g}$ $s_{(R)} = \pm 0.037 \text{ g}/100 \text{ g}$

D-fructose:

$x = 6.72 \text{ g}/100 \text{ g}$ $r = 0.587 \text{ g}/100 \text{ g}$ $s_{(r)} = \pm 0.207 \text{ g}/100 \text{ g}$
 $R = 0.748 \text{ g}/100 \text{ g}$ $s_{(R)} = \pm 0.264 \text{ g}/100 \text{ g}$

sucrose:

$x = 43.32 \text{ g}/100 \text{ g}$ $r = 1.722 \text{ g}/100 \text{ g}$ $s_{(r)} = \pm 1.033 \text{ g}/100 \text{ g}$
 $R = 4.268 \text{ g}/100 \text{ g}$ $s_{(R)} = \pm 1.501 \text{ g}/100 \text{ g}$

For further data see references (Ref. C 2.4)

6. Recognizing interference during the assay procedure

6.1 If the conversion of D-glucose has been completed according to the time given under "Procedure", it can be concluded in general that no interference has occurred.

6.2 On completion of the reaction, the determination can be restarted by adding D-glucose (qualitative or quantitative): if the absorbance is altered subsequent to the addition of the standard material, this is also an indication that no interference has occurred.

6.3 Operator error or interference of the determination through the presence of substances contained in the sample can be recognized by carrying out a double determination using two different sample volumes (e.g. 0.100 ml and 0.200 ml): the measured differences in absorbance should be proportional to the sample volumes used.

When analyzing solid samples, it is recommended that different quantities (e.g. 1 g and 2 g) be weighed into 100 ml volumetric flasks. The absorbance differences measured and the weights of sample used should be proportional for identical sample volumes.

6.4 Possible interference caused by substances contained in the sample can be recognized by using an internal standard as a control: in addition to the sample, blank and standard determinations, a further determination should be carried out with sample and assay control solution in the same assay. The recovery can then be calculated from the absorbance differences measured.

6.5 Possible losses during the determination can be recognized by carrying out recovery tests: the sample should be prepared and analyzed with and without added standard material. The additive should be recovered quantitatively within the error range of the method.

7. Reagent hazard

The reagents used in the determination of D-glucose are not hazardous materials in the sense of the Hazardous Substances Regulations, the Chemicals Law or EC Regulation 67/548/EEC and subsequent alteration, supplementation and adaptation guidelines. However, the general safety measures that apply to all chemical substances should be adhered to.

After use, the reagents can be disposed of with laboratory waste, but local regulations must always be observed. Packaging material can be disposed of in waste destined for recycling.

8. General information on sample preparation

In carrying out the assay:

Use **clear, colorless and practically neutral liquid samples** directly, or after dilution according to the dilution table, and of a volume up to 2.000 ml; Filter **turbid solutions**;

Degas **samples containing carbon dioxide** (e.g. by filtration);

Adjust **acid samples** to approx. pH 8 by adding sodium or potassium hydroxide solution;

Adjust **acid and weakly colored samples** to approx. pH 8 by adding sodium or potassium hydroxide solution and incubate for approx. 15 min; Measure **"colored" samples** (if necessary adjusted to approx. pH 8) against a sample blank (= buffer or redist. water + sample), adjust the photometer to 0.000 with the blank in the beam;

Treat **"strongly colored" samples** that are used undiluted or with a higher sample volume with polyvinylpyrrolidone (PVPP) or with polyamide, e.g. 1 g/100 ml;

Crush or homogenize **solid or semi-solid samples**, extract with water or dissolve in water and filter if necessary; resp. remove turbidities or dyestuffs by Carrez clarification;

Deproteinize **samples containing protein** with Carrez reagents;

Extract **samples containing fat** with hot water (extraction temperature should be above the melting point of the fat involved). Cool to allow the fat to separate, make up to the mark, place the volumetric flask in an ice bath for 15 min and filter; alternatively clarify with Carrez-solutions after the extraction with hot water.

Carrez clarification:

Pipette the liquid sample into a 100 ml volumetric flask which contains approx. 60 ml redist. water, or weigh sufficient quantity of the sample into a 100 ml volumetric flask and add approx. 60 ml redist. water. Subsequently, carefully add 5 ml Carrez-I-solution (potassium hexacyanoferrate(II) (ferrocyanide), 85 mM = 3.60 g $\text{K}_4[\text{Fe}(\text{CN})_6] \times 3 \text{ H}_2\text{O}/100 \text{ ml}$) and 5 ml Carrez-II-solution (zinc sulfate, 250 mM = 7.20 g $\text{ZnSO}_4 \times 7 \text{ H}_2\text{O}/100 \text{ ml}$). Adjust to pH 7.5-8.5 with sodium hydroxide (0.1 M; e.g. 10 ml). Mix after each addition. Fill the volumetric flask to the mark, mix and filter.

Samples containing protein should only be deproteinized with perchloric acid or with trichloroacetic acid in the absence of sucrose and maltose as these disaccharides are fully or partially hydrolyzed with the release of D-glucose. The Carrez clarification is recommended for normal use.

9. Application examples

Determination of D-glucose in milk

Pipette 20 ml milk into a 100 ml volumetric flask, add the following solutions and mix after each addition: 10 ml Carrez-I-solution (see pt. 8), 10 ml Carrez-II-solution (see pt. 8) and 20 ml sodium hydroxide (0.1 M). Fill up to the mark with redist. water, mix and filter. Use 1.000 ml or 2.000 ml, respectively, of the filtrate for the assay.

Determination of carbohydrates in diet beer for diabetics (total D-glucose after acid hydrolysis) (Ref. A 2.4)

Shake 300 to 500 ml of beer in order to remove carbon dioxide and filter through fluted filter paper. Pipette 50 ml filtrate into a 250 ml flask, add 10 ml HCl (25%; $d = 1.125$) and 50 ml redist. water, boil for 3 h under reflux in a bath with boiling water. Cool down to 20-25°C, adjust the pH 6-7 by the addition of NaOH (5 M), transfer into a 250 ml volumetric flask, fill up to the mark with redist. water, mix and filter. Use the 1:10 (1+9) with redist. water diluted filtrate for the assay.

Determination of D-glucose in baking agents, spices and salt mixtures

Accurately weigh the homogenized sample into a volumetric flask and dissolve with water, resp. extract; if necessary, incubate for 30 min at 60-70°C. Remove turbidities by means of Carrez reagents. Use the sample solution, after dilution according to the dilution table, for the assay.

10. Further applications

The method may also be used in the examination of cosmetics (Ref. B 3.10), pharmaceuticals (Ref. B 3.6), paper (Ref. D 2.2), and tobacco (Ref. C 3.7) and in research when analyzing biological samples. For sampling, treatment and stability of the sample see Ref. A 1.1, A 1.2.

Determination of D-glucose in fermentation samples and cell culture media

Place the sample (after centrifugation, if necessary) in a water-bath at 80°C for 15 min to stop enzymatic reactions. Centrifuge and use the supernatant (diluted according to the dilution table, if necessary) for the assay. Alternatively, deproteinization can be carried out with perchloric acid, but only by absence of disaccharides, or with Carrez-solutions. See the above-mentioned examples.

Homogenize gelatinous agar media with water and treat further as described.

References

A. References for the determination of D-glucose

- A 1.1 Bergmeyer, H.U., Bernt, E., Schmidt, F. & Stork, H. (1974) in *Methoden der enzymatischen Analyse* (Bergmeyer, H.U., Hrsg.) 3. Aufl., Bd. 2, 1241-1246; Verlag Chemie, Weinheim, and (1974) in *Methods of Enzymatic Analysis* (Bergmeyer, H.U., ed.) 2nd ed., vol. 3, pp. 1196-1201; Verlag Chemie, Weinheim/Academic Press, Inc., New York and London
- A 1.2 Kunst, A., Draeger, B. & Ziegenhorn, J. (1984) in *Methods of Enzymatic Analysis* (Bergmeyer, H.U., Hrsg.) 3rd ed., vol. VI, pp. 163-172, Verlag Chemie, Weinheim, Deerfield Beach/Florida, Basel
- A 2.1 Brautechnische Analysenmethoden, Band II, S. 400-402 (1979), Methodensammlung der Mitteleuropäischen Brautechnischen Analytikkommission (MEBAK), herausgegeben von F. Drawert im Selbstverlag der MEBAK, Freising
- A 2.2 Schweizerisches Lebensmittelbuch (1981), Kapitel 61 B (Enzymatische Bestimmungen)/1.1
- A 2.3 Gombocz, E., Hellwig, E., Vojir, F. & Petuely, F. (1981) *Deutsche Lebensmittel-Rundschau* **77**, 3 (Glucose)
- A 2.4 Amtliche Sammlung von Untersuchungsverfahren nach §35 LMBG; Untersuchung von Lebensmitteln: Bestimmung von Glucose in Fleischerzeugnissen, 07.00-22 (Mai 1983); Bestimmung von Glucose in Würstwaren, 08.00-23 (Mai 1983); Bestimmung der belastenden Kohlenhydrate in Diätbier für Diabetiker (als Gesamtglucose), 49.01.05-1 (Mai 1984)
- A 2.5 Ministero dell'Agricoltura e delle Foreste (1986) Approvazione dei "Metodi ufficiali di analisi per i mosti, i vini, gli agri di vino (aceti) e i sottoprodotti della vinificazione". Gazzetta Ufficiale della Repubblica Italiana, n. 161 del 14 luglio 1986
- A 3.1 Henniger, G. & Schütz, A. (1987) *Methods for the Enzymatic Determination of D-Glucose*, Poster-Präsentation bei Association of Official Analytical Chemists Annual International Meeting, San Francisco, CA, USA

For further information see instructions for

Test-Combination D-Glucose/D-Fructose	Cat. No. 10 139 106 035
Test-Combination Maltose/Sucrose/ D-Glucose	Cat. No. 11 113 950 035
Test-Combination Sucrose/D-Glucose	Cat. No. 10 139 041 035
Test-Combination Sucrose/D-Glucose/ D-Fructose	Cat. No. 10 716 260 035
Test-Combination D-Sorbitol/Xylitol	Cat. No. 10 670 057 035
Test-Combination Starch	Cat. No. 10 207 748 035

B. References for the determination of D-glucose and D-fructose

- B 1.1 Schmidt, F.H. (1961) Die enzymatische Bestimmung von Glucose und Fructose nebeneinander, *Klinische Wochenschrift* **39**, 1244-1247
- B 1.2 Bernt, E. & Bergmeyer, H.U. (1974) in *Methoden der enzymatischen Analyse* (Bergmeyer, H.U., Hrsg.) 3. Aufl., Bd. 2, S. 1349-1352; Verlag Chemie, Weinheim and (1974) in *Methods of Enzymatic Analysis* (Bergmeyer, H.U., ed.) 2nd ed., vol. 3, pp. 1304-1307; Verlag Chemie, Weinheim/Academic Press, Inc., New York and London
- B 1.3 Beutler, H.O. (1984) in *Methods of Enzymatic Analysis* (Bergmeyer, H.U., ed.) 3rd ed., vol. VI, pp. 321-327, Verlag Chemie, Weinheim, Deerfield Beach/Florida, Basel
- B 2.1 Handbuch für diätetische Lebensmittel. Analytik - Reinheitsanforderungen - (1973) des Bundesverbandes der diätetischen Lebensmittelindustrie e.V., Bad Homburg, herausgegeben von der Fördergesellschaft Diätetische Lebensmittel mbH, S. 27-30
- B 2.2 Lafon-Lafourcade, S., Lafitte, M. & Joyeux, A. (1977) Dosage du Glucose et du Fructose Residuels dans les Vins par Methode Enzymatique, Office International de la Vigne et du Vin, n 600/F.V. 634
- B 2.3 Deutsche Norm DIN 10381 (April 1979) Untersuchung von Stärke und Stärkeerzeugnissen; Bestimmung von D-Glucose und D-Fructose in derselben Untersuchungssprobe (Enzymatisches Verfahren),
- B 2.4 Methodenbuch für Weinanalysen in Österreich (1980), herausgegeben von Arbeitsgemeinschaft der Landw. Versuchsanstalten in Österreich (ALVA)
- B 2.5 Schweizerisches Lebensmittelbuch, Kapitel 61B (Enzymatische Bestimmungen)/1.1, 1.2 und 1.6 (1981), Kapitel 2A (Milchmischgetränke)/11 (1980), Kapitel 9 (Speiseeis)/ 4.3 (1983), Kapitel 22 (Diätetische Lebensmittel und Speziallebensmittel)/6.3 (1991) Kapitel 23A (Honig)/8.2 (1995), Kapitel 28A (Frucht- und Gemüsesäfte u.a.)/5.4 (1988), Kapitel 30A (Wein aus Trauben)/4.4 (1993), Kapitel 34 (Gärungssessig)/8.1 (1994)
- B 2.6 Gombocz, E., Hellwig, E., Vojir, F. & Petuely, F. (1981) *Deutsche Lebensmittel-Rundschau* **77**, 3 (Glucose) und 9-10 (Fructose)
- B 2.7 Brautechnische Analysenmethoden, Band III, S. 580-586 (1982), Methodensammlung der Mitteleuropäischen Brautechnischen Analytikkommission (MEBAK), herausgegeben von F. Drawert im Selbstverlag der MEBAK, Freising
- B 2.8 Österreichisches Lebensmittelbuch (Codex Alimentarius Austriacus) Kapitel B15 (Kakao, Kakaoverzeugnisse, Lebensmittel mit Kakao oder Schokolade, Nougat, Nougatmassen), 1983; Kapitel B22 (Zucker und Zuckerarten) (1983)
- B 2.9 Amtliche Sammlung von Untersuchungsverfahren nach §35 LMBG; Untersuchung von Lebensmitteln: Bestimmung von Glucose und Fructose in Fruchtsäften, 31.00-12 (November 1984); Bestimmung von Glucose und Fructose in Kinder-Zwieback und Zwiebackmehl, 48.02.07-1 (Mai 1985); Enzymatische Bestimmung der Gehalte an D-Glucose und D-Fructose in Frucht- und Gemüsesäften, 31.00-12 (Januar 1997); Enzymatische Bestimmung der Gehalte an D-Glucose und D-Fructose in Gemüsesäften, 26.26-11 (Januar 1997);
- B 2.10 Henniger, G. & Mascaro, L. (1985) Enzymatic-Ultraviolet Determination of Glucose and Fructose in Wine: Coll. Study, J. Assoc. Off. Anal. Chem. **68**, 1021-1024
- B 2.11 Official Methods of Analysis of the Association of Official Analytical Chemists (1990), 15th ed., vol. 2, pp. 741-742 (985.09)
- B 2.12 The method is approved for official wine inspection in Rhenish-Palatinate (1985; Chamber of Agriculture, Bad Kreuznach) and in Hesse (1986, Ministry for Agriculture and Forestry).
- B 2.13 Ministero dell'Agricoltura e delle Foreste (1986) Approvazione dei "Metodi ufficiali di analisi per i mosti, i vini, gli agri di vino (aceti) e i sottoprodotti della vinificazione". Gazzetta Ufficiale della Repubblica Italiana, n. 161 del 14 luglio 1986
- B 2.14 International Federation of Fruit Juice Producers (IFU, Methods of Analysis, no. 55-1985); contained in "Code of Practice for Evaluation of Fruit and Vegetable Juices" (1996) edited by Association of the Industry of Juices and Nectars from Fruits and Vegetables of the European Economic Community (A.I.J.N.)
- B 2.15 RSK-Values, The Complete Manual, Guide Values and Ranges of Specific Numbers for Fruit Juices and Nectars, Including the Revised Methods of Analysis (1987), 1st ed., Verlag Flüssiges Obst/Liquid Fruit, D-56370 Eschborn, pp. 97-100
- B 2.16 Nederlandse Norm NEN 2857 (1e druk, oktober 1989) Vruchtesappen: Bepaling van het glucose- en fructosegehalte; Enzymatische methode (Fruit juices - Determination of the glucose and fructose content - Enzymatic method)
- B 2.17 Recueil des méthodes internationales d'analyse des vins et des moûts, Complément n° 1 à l'édition officielle de juin 1990, OFFICE INTERNATIONAL DE LA VIGNE ET DU VIN, S. 97-100
- B 2.18 Official Journal of the European Communities L 272 (3 October 1990), Legislation: Commission Regulation (EEC) No 2676/90 of 17 September 1990 determining Community methods for the analysis of wines (pp. 61-63)
- B 2.19 Deutsche Norm DIN EN 1140 (1994) Frucht- und Gemüsesäfte; Enzymatische Bestimmung der Gehalte an D-Glucose und D-Fructose; Spektralphotometrische Bestimmung von NADPH (Fruit and vegetable juices; Enzymatic determination of D-glucose and D-fructose content; NADPH spectrophotometric method)
- B 2.20 European Standard EN 1140 (Dec. 1994) Fruit and vegetable juices; Enzymatic determination of D-glucose and D-fructose content by the NADPH spectrometric method
- B 2.21 Standard der Russischen Föderation / Standard of the Russian Federation / GOSSTANDART ROSSII GOST R 51240-98 (1998) Fruit and vegetable juices. Determination of D-glucose and D-fructose content
- B 3.1 Weichel, H.H. (1965) Die quantitative Bestimmung von Fructose neben anderen Kohlenhydraten in Lebensmitteln durch enzymatische Analyse, *Deutsche Lebensmittel-Rundschau* **61**, 53-55
- B 3.2 Tschersich, J. & Mauch, W. (1968) Enzymatisch-photometrische Bestimmung von D-Glucose und D-Fructose in Verbraucherzucker, *Zeitschrift für die Zuckerindustrie* **18**, 107-110
- B 3.3 Baumann, G. & Gierschner, K. (1971) Die Bestimmung von Zuckern in Fruchtsäften - ein Vergleich der enzymatischen mit der Luff-Schoorl-Methode, *Ind. Obst- und Gemüseverwertung* **56**, 165-170
- B 3.4 Kubadinow, N. (1974) Die enzymatische Bestimmung von D-Glucose und D-Fructose in Zuckerrüben und Betriebsäften der Zuckerproduktion und Vergleich ihrer Ergebnisse mit den Werten der Berliner Institutsmethode, *Zucker* **27**, 72-78
- B 3.5 Schiweck, H. & Büsching, L. (1974) Das Verhalten von Glucose und Fructose während der Zuckerfabrikation, *Zucker* **27**, 122-128
- B 3.6 Schultze, K.W. & Rummel, M. (1975) Beispiel für die Anwendung von Enzymen bei der pharmazeutischen Analyse, *Acta Pharmaceutica Technologica* **21**, 167-170

- B 3.7 Promayon, J., Barel, M., Fourny, G. & Vincent, J.-C. (1976) Essais de Détermination de la Teneur en Chicorée dans les Mélanges Solubles de Café et de Chicorée, *Café Cacao Thé* **20**, 209-217
- B 3.8 Motz, R.J. (1976) Enzymatische Zuckerbestimmung, *Zucker und Süßwarenwirtschaft* **29**, 75-77
- B 3.9 Wagner, K. & Kreutzer, P. (1977) Zusammensetzung und Beurteilung von Auslesen, Beeren- und Trockenbeerenauslesen, *Die Weinwirtschaft* **10**, 272-275
- B 3.10 Henniger, G. & Boos, H. (1978) Anwendung der enzymatischen Analyse bei der Untersuchung kosmetischer Präparate - dargestellt an einigen Beispielen, Seifen - Öle - Fette - Wachse **104**, 159-164
- B 3.11 Klingebiel, L., Großklaus, R. & Pahlke, G. (1979) Erfahrungen mit der enzymatischen Bestimmung von Zucker und Zuckeraustauschstoffen in Diabetikerwaren, *Z. Lebensm. Unters. Forsch.* **169**, 359-360
- B 3.12 Wucherpfennig, K., Otto, K. & Yueh-Chyng Huang (1986) Aussagekraft des Glucose-Fructose-Verhältnisses, *Die Weinwirtschaft-Technik* **122**, 254-260
- B 3.13 Millies, K. D., Sponholz, W.R. & Eberle, C. (1988) Methoden zur Zuckerbestimmung, *Weinwirtschaft - Technik*, Heft 6, S. 6-11
- B 3.14 Plessi, M., Monzani, A., & Coppini, D. (1988) Determination of the Monosaccharide and Alcohol Content of Balsamic and Other Vinegars by Enzymatic Methods, *Agric.Biol.Chem.* **52**, 25-30

C. References for the determination of sucrose, D-glucose and D-fructose

- C 2.1 Fuchs, G. & Wretling S. (1979) Bestämning av fruktos, glukos och sackaros i livsmedel, *Vår Föda* **31**, 435-439
- C 2.2 Bundesverband der Deutschen Feinkostindustrie e.V. Bonn; Analysemethoden: Bestimmung des Zuckergehaltes in Tomatenmark (enz.), IV/61 (Dezember 1979)
- C 2.3 Norme Française Homologuée NF V 76-106 (Octobre 1980) Jus Fruits et Jus de Légumes: Détermination de la Teneur en Saccharose, D-Glucose, D-Fructose,
- C 2.4 Amtliche Sammlung von Untersuchungsverfahren nach § 35 LMBG; Untersuchung von Lebensmitteln: Bestimmung des Zuckergehalts in Tomatenmark, 26.11.03-8 (Mai 1983); Bestimmung des Zuckergehalts in Tomatenketchup und vergleichbaren Erzeugnissen, 52.01.01-8 (November 1983); Bestimmung von Saccharose, Glucose und Fructose in teildaptierter Säuglingsnahrung auf Milchbasis, 48.01-3 (Mai 1985); Bestimmung des Gesamtzuckergehaltes in Speisesenf, 52. 06-5 (Dezember 1991); Bestimmung von Glucose, Fructose und Saccharose in Eiern und Eiprodukten, 05.00-10 (Juni 1992)
- C 2.5 Office International du Cacao, du Chocolat et de la Confiserie, IOCCC Method number 113-1989: Determination of Glucose, Fructose and Sucrose in Chocolate and Sugar Confectionery Products by Means of Enzymes, Draft Standard Method
- C 2.6 Österreichisches Lebensmittelbuch (Codex Alimentarius Austriacus), Kapitel B29: Senf; Erlaß vom 27. Oktober 1989 (s. ERNÄHRUNG/NUTRITION **14**, 168-170 (1990))
- C 3.1 Drawert, F. (1964) Enzymatische Analyse von Glucose, Fructose, Saccharose und Sorbit in Weinen und Traubenmosten, *VITIS* **4**, 185-187
- C 3.2 Trautner, K. (1969) Enzymatische Zuckerbestimmungen, *Zeitschrift für Ernährungswissenschaft, Suppl.* **8**, S. 40-45
- C 3.3 Somogyi, J.C. & Trautner, K. (1974) Der Glucose-, Fructose- und Saccharosegehalt verschiedener Gemüsearten, *Schweiz. Med. Wochenschrift* **104**, 177-182
- C 3.4 Trautner, K. & Somogyi, J.C. (1979) Zuckergehalte von Obst und Gemüse - Einflüsse von Reifegrad, Sorte und Lagerung, *Mitt. Gebiete Lebensm. Hyg.* **67**, 497-508
- C 3.5 Zürcher, K. & Hadorn, H. (1976) Veränderungen des Zuckerspektrums eines Sirups während der Lagerung, *Mitt.Gebiete Lebensm.Hyg.* **72**, 136-139
- C 3.6 Zürcher, K. & Hadorn, H. (1976) Vergleichende Zuckerbestimmungen mit gaschromatographischen, enzymatischen und reduktometrischen Methoden, *Deutsche Lebensmittel-Rundschau* **72**, 197-202
- C 3.7 Sekin, S. (1978) Enzymatic Determination of Glucose, Fructose and Sucrose in Tobacco, *Tobacco Sci.* , 75-77; *Tobacco International* **181**, 27-29
- C 3.8 Polascsek-Rác, M., Pauli, M. P., Horváth, G. & Vámos-Vigyázó (1981) Enzymatic determination of the sugars in red pepper, *Z. Lebensm. Unters. Forsch.* **172** , 115-117

D. References for the determination of sucrose and D-glucose

- D 1.1 Bergmeyer, H. U. & Bernt, E. (1974) in *Methoden der enzymatischen Analyse* (Bergmeyer, H.U., Hrsg.) 3. Aufl., Bd. 2, S. 1221-1224; Verlag Chemie, Weinheim, and (1974) in *Methods of Enzymatic Analysis* (Bergmeyer, H. U., ed.) 2nd ed., vol. 3, p. 1176-1179, Verlag Chemie Weinheim/Academic Press, Inc., New York and London
- D 2.1 *Handbuch für diätetische Lebensmittel. Analytik - Reinheitsanforderungen -* (1973) des Bundesverbandes der diätetischen Lebensmittelindustrie e.V., Bad Homburg, herausgegeben von der Fördergesellschaft Diätetische Lebensmittel mbH, S. 57-60
- D 2.2 *Untersuchung von Papieren, Kartons und Pappen für Lebensmittelverpackungen* (gem. Empfehlungen XXXVI der Kunststoffkommission des Bundesgesundheitsamtes) Kapitel 8 (Methoden), Pkt. 3.5.2 (März 1979)
- D 2.3 Schweiz. Lebensmittelbuch, Kapitel 61B (Enzymatische Bestimmungen)/1.3 (1981), Kap. 2A (Milchmischgetränke)/10 (1980), Kap. 2B (Sauermilchprodukte)/10 (1980), Kapitel 4 (Milchdauerwaren)/5.2 (1993), Kap. 9 (Speiseeis)/4.3 (1983), Kap. 22 (Diätetische Lebensmittel und Speziallebensmittel)/6.3 (1991), Kap. 28A (Frucht- und Gemüsesäfte u.a.)/5.4 (1988), Kapitel 30A (Wein aus Trauben)/4.4 (1993), Kap. 36A (Kakao, Kakaomasse, Kakaopulver und Schokoladenpulver)/7.2 (1992)
- D 2.4 Gombocz, E., Hellwig, E., Vojir, F. & Petuely, F. (1981) *Deutsche Lebensmittel-Rundschau* **77**, 3 (Glucose), 11 (Saccharose)
- D 2.5 Brautechnische Analysemethoden, Band III, S. 586-589 (1982), *Methodensammlung der Mitteleuropäischen Brautechnischen Analytikkommission (MEBAK)*, herausgegeben von F. Drawert im Selbstverlag der MEBAK, Freising
- D 2.6 Amtliche Sammlung von Untersuchungsverfahren nach §35 LMBG; Untersuchung von Lebensmitteln: Bestimmung von Saccharose in Fleischerzeugnissen, 07.00-24, (Mai 1983); Bestimmung von Saccharose in Wurstwaren, 08.00-25 (Mai 1983); Bestimmung von Saccharose in Fruchtsäften, 31.00-13 (November 1984); Bestimmung des Gehaltes an Saccharose und Glucose in Milchprodukten und Speiseeis, 02.00-12 (Mai 1986); Bestimmung des Gehaltes an Saccharose und Glucose in Käse, 03.00-12 (Mai 1986); Bestimmung des Gehaltes an Saccharose und Glucose in Speiseeis, 42.00-5 (Mai 1986); Enzymatische Bestimmung des Saccharosegehaltes in Frucht- und Gemüsesäften, 31.00-13 (September 1997); Enzymatische Bestimmung des Saccharosegehaltes in Gemüsesäften, 26.26-17 (September 1997)
- D 2.7 Deutsche Norm DIN 10326 (Februar 1986) Bestimmung des Gehaltes an Saccharose und Glucose in Milchprodukten und Speiseeis
- D 2.8 International Federation of Fruit Juice Producers (IFU, Methods of Analysis, no. 56-1985); contained in "Code of Practice for Evaluation of Fruit and Vegetable Juices" (1996) edited by Association of the Industry of Juices and Nectars from Fruits and Vegetables of the European Economic Community (A.I.J.N.)
- D 2.9 Österreichisches Lebensmittelbuch (Codex Alimentarius Austriacus), Kapitel B8 (Essig), Erlaß vom 16. Juni 1986 (s. ERNÄHRUNG/NUTRITION **11**, 49-53 (1987); Kapitel B15 (Kakao, Kakaoverzeugnisse, Lebensmittel mit Kakao oder Schokolade, Nougat, Nougatmassen) (1983)
- D 2.10 Niederlande: Warenwet, Uitvoeringsvoorschriften (CII-6) Regeling Onderzoekingsmethoden voor brood; Methode 14: De Bepaling van het Suikergehalte als de Som van Saccharose en Invertsuiker, de laatste herleid tot Saccharose (Oktober 1986); Dit voorschrift betreft een methode voor de enz. bepaling van de som van saccharose en invertsuiker, de laatste berekend als saccharose, in brood
- D 2.11 RSK-Values, The Complete Manual, Guide Values and Ranges of Specific Numbers for Fruit Juices and Nectars, Including the Revised Methods of Analysis (1987), 1st ed., Verlag Flüssiges Obst/Liquid Fruit, D-56370 Eschborn, pp. 145-148
- D 2.12 Verband Deutscher Landwirtschaftlicher Untersuchungs- und Forschungsanstalten, VDLUFA (1988) *Methodenbuch Band VI: Enzymatische Bestimmung des Glucose- und Saccharosegehaltes von Milchprodukten (C20.3)*
- D 2.13 Niederlande Norm NEN 2858 (1e druk, oktober 1989) Vruchtesappen: Bepaling van het saccharosegehalte; Enzymatische methode (Fruit juices - Determination of the sucrose content - Enzymatic method)
- D 2.14 Europäische Norm/European Standard EN 12146 (Okt. 1996) Frucht- und Gemüsesäfte - Enzymatische Bestimmung des Saccharosegehaltes - Spektralphotometrisches Verfahren mit NADP (Fruit and vegetable juices - Enzymatic determination of sucrose content - NADP spectrophotometric method)
- D 2.15 Deutsche Norm DIN EN 12146 (Okt. 1996) Frucht- und Gemüsesäfte, Teil 4: Enzymatische Bestimmung des Saccharosegehaltes; Spektralphotometrisches Verfahren mit NADP.
- D 2.16 Niederlande Norm NEN-EN 12146 (Oktober 1996) Vruchten- en groentesappen. Enzymatische Bepaling van het gehalte aan saccharose. NADP-spectrometrische methode
- D 2.17 Standard der Russischen Föderation / Standard of the Russian Federation / GOSSTANDART ROSSII GOST R 51258-99 (1999) Milk and milk products. Method for determination of sucrose and glucose content

D-Glucose assay control solution (Bottle 3)

Concentration*: see bottle label

D-Glucose assay control solution is a stabilized aqueous solution of D-glucose. It serves as assay control solution for the enzymatic analysis of D-glucose in foodstuffs and other materials.

Application:

1. *Addition of D-glucose assay control solution to the assay mixture:*
Instead of sample solution the assay control solution is used for the assay.

2. *Restart of the reaction, quantitatively:*

After completion of the reaction with sample solution and measuring of A_2 , add 0.050 ml assay control solution to the assay mixture. Read absorbance A_3 after the end of the reaction (approx. 15 min.). Calculate the concentration from the difference of ($A_3 - A_2$) according to the general equation for calculating the concentration. The altered total volume must be taken into account. Because of the dilution of the assay mixture by addition of the assay control solution, the result differs insignificantly from the data stated on the bottle label.

3. Internal standard:

The assay control solution can be used as an internal standard in order to check the determination for correct performance (gross errors) and to see whether the sample solution is free from interfering substances:

Pipette into cuvettes	Blank	Sample	Standard	Sample + Standard
solution 1	1.000 ml	1.000 ml	1.000 ml	1.000 ml
sample solution	-	0.100 ml	-	0.050 ml
assay control sln.	-	-	0.100 ml	0.050 ml
redist. water	2.000 ml	1.900 ml	1.900 ml	1.900 ml

Mix, and read absorbances of the solutions (A_1) after approx. 3 min. Continue as described in the pipetting scheme under "Procedure". Follow the instructions given under "Instructions for performance of assay" and the footnotes.

The recovery of the standard is calculated according to the following formula:

$$\text{recovery} = \frac{2 \times \Delta A_{\text{sample + standard}} - \Delta A_{\text{sample}}}{\Delta A_{\text{standard}}} \times 100 [\%]$$

* Stated as anhydrous D-glucose

